



# Care pathway for hypertension



<sup>3</sup> Signs of papilloedema or retinal haemorrhage.  
<sup>4</sup> Labile or postural hypotension, headache, palpitations, pallor and diaphoresis.  
<sup>5</sup> Ambulatory blood pressure monitoring.  
<sup>6</sup> Home blood pressure monitoring.

## Summary of antihypertensive drug treatment

Also see 'Initiating and titrating antihypertensive drug treatment' on pages 11–12.



<sup>18</sup> Choose a low-cost ARB.

<sup>19</sup> A CCB is preferred but consider a thiazide-like diuretic if a CCB is not tolerated or the person has oedema, evidence of heart failure or a high risk of heart failure.

<sup>20</sup> Consider a low dose of spironolactone<sup>21</sup> or higher doses of a thiazide-like diuretic.

<sup>21</sup> At the time of publication (August 2011), spironolactone did not have a UK marketing authorisation for this indication. Informed consent should be obtained and documented.

<sup>22</sup> Consider an alpha-blocker or beta-blocker if further diuretic therapy is not tolerated, or is contraindicated or ineffective.

## Key priorities for implementation *continued*

### Choosing antihypertensive drug treatment

- Offer people aged 80 years and over the same antihypertensive drug treatment as people aged 55–80 years, taking into account any comorbidities.

#### Step 1 treatment

- Offer step 1 antihypertensive treatment with a calcium-channel blocker (CCB) to people aged over 55 years and to black people of African or Caribbean family origin of any age. If a CCB is not suitable, for example because of oedema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.
- If a diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as chlortalidone (12.5–25.0 mg once daily) or indapamide (1.5 mg modified-release or 2.5 mg once daily) in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.
- For people who are already having treatment with bendroflumethiazide or hydrochlorothiazide and whose blood pressure is stable and well controlled, continue treatment with the bendroflumethiazide or hydrochlorothiazide.

#### Step 4 treatment

- For treatment of resistant hypertension at step 4:
  - Consider further diuretic therapy with low-dose spironolactone (25 mg once daily)<sup>2</sup> if the blood potassium level is 4.5 mmol/l or lower. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalaemia.
  - Consider higher-dose thiazide-like diuretic treatment if the blood potassium level is higher than 4.5 mmol/l.

<sup>2</sup> At the time of publication (August 2011), spironolactone did not have a UK marketing authorisation for this indication. Informed consent should be obtained and documented.

## Monitoring treatment

- Use clinic blood pressure measurement to monitor the response to treatment.
- For people identified as having a 'white-coat effect'<sup>23</sup>, consider ABPM or HBPM as an adjunct to clinic blood pressure measurements to monitor the response to treatment.

### Blood pressure targets

#### Clinic blood pressure

- People aged under 80 years: lower than 140/90 mmHg
- People aged over 80 years: lower than 150/90 mmHg

#### Daytime average ABPM or average HBPM blood pressure during the person's usual waking hours

- People aged under 80 years: lower than 135/85 mmHg
- People aged over 80 years: lower than 145/85 mmHg

## Patient education and adherence to treatment

- Help people to make informed choices by providing guidance and materials about the benefits of drugs and the unwanted side effects sometimes experienced<sup>24</sup>.
- Tell people about patient organisations that have forums for sharing views and information<sup>24</sup>.
- Offer an annual review of care to monitor blood pressure, provide people with support and discuss their lifestyle, symptoms and medication<sup>24</sup>.

### Interventions to support adherence to treatment

- Only use interventions to overcome practical problems associated with non-adherence if a specific need is identified<sup>25</sup>.
- Target the intervention to the need. Interventions might include:
  - suggesting that people record their medicine-taking
  - encouraging people to monitor their condition
  - simplifying the dosing regimen
  - using alternative packaging for the medicine
  - using a multi-compartment medicines system<sup>25</sup>.

<sup>23</sup> A discrepancy of more than 20/10 mmHg between clinic and average daytime ABPM or average HBPM measurements at the time of diagnosis.

<sup>24</sup> This recommendation was developed for the original 2004 guideline.

<sup>25</sup> This recommendation is taken from 'Medicines adherence' (NICE clinical guideline 76, 2009), available from [www.nice.org.uk/guidance/CG76](http://www.nice.org.uk/guidance/CG76)

## Diagnosing hypertension

### Measuring the clinic blood pressure

- Measure blood pressure in both arms.
  - If the difference in readings between arms is more than 20 mmHg, repeat the measurements.
  - If the difference in readings between arms remains more than 20 mmHg on the second measurement, measure subsequent blood pressures in the arm with the higher reading.
- If blood pressure measured in the clinic is 140/90 mmHg or higher:
  - Take a second measurement during the consultation.
  - If the second measurement is substantially different from the first, take a third measurement.

Record the lower of the last two measurements as the clinic blood pressure.

### Confirming the diagnosis

- If the clinic blood pressure is 140/90 mmHg or higher, offer ABPM to confirm the diagnosis of hypertension.
- If a person is unable to tolerate ABPM, HBPM is a suitable alternative to confirm the diagnosis of hypertension.
- While waiting to confirm the diagnosis, carry out investigations for target organ damage and a formal assessment of cardiovascular risk (see page 10).

### Severe hypertension

- Consider starting antihypertensive drug treatment immediately, without waiting for the results of ABPM or HBPM, for people with severe hypertension.

### Specialist investigations

- Refer people to specialist care the same day if they have:
  - accelerated hypertension (blood pressure usually higher than 180/110 mmHg with signs of papilloedema and/or retinal haemorrhage) **or**
  - suspected pheochromocytoma (labile or postural hypotension, headache, palpitations, pallor and diaphoresis).
- Consider the need for specialist investigations in people with signs and symptoms suggesting a secondary cause of hypertension.